

HD-A137 968

MOMENTUM DEPENDENT LINE SHAPES IN PHOTOEMISSION(U)
PENNSYLVANIA UNIV PHILADELPHIA R A DIDIO ET AL.
17 JAN 84 TR-22 N00014-79-C-0991


1/1

UNCLASSIFIED

F/G 20/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 22	2. GOVT ACCESSION NO. AD-A137968	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Momentum Dependent Line Shapes in Photoemission		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) R.A. DiDio and E.W. Plummer, Dept. of Physics, W.R. Graham, LRSM, University of Pennsylvania, Philadelphia, PA 19104		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0991
9. PERFORMING ORGANIZATION NAME AND ADDRESS Trustees of the Univ. of Pennsylvania Office of Projects Research and Grants 3451 Walnut St. Philadelphia, PA 19104		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR SRO-013/9-12-79 (472)
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Dept. of the Navy 500 N. Quincy St. Arlington, VA 22217		12. REPORT DATE Jan. 17, 1984
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 14
		15. SECURITY CLASS. (of this report) Unclassified
		16. DECLASSIFICATION SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DTIC S ELECT FEB 16 1984		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) line shapes, valence levels, angle resolved photoelectron spectroscopy		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The crystal-momentum dependence of the line shapes appearing in photo-excitation from the valence levels of ordered S on Fe(100) has been measured using angle resolved photoelectron spectroscopy. The variation in line shape with k is a consequence of the hybridization of the energy bands of the 2-D overlayer with the 3-D solid. This observation, predicted theoretically by Liebsch, yields detailed information about the bonding interactions of ordered chemisorption systems.		

AD A137968

DTIC FILE COPY

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6001

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

84 02 16 132

Momentum Dependent Line Shapes in Photoemission

R.A. DiDio and E.W. Plummer

Department of Physics and

Laboratory for Research on the Structure of Matter

University of Pennsylvania

Philadelphia, PA 19104

and

W.R. Graham

Department of Materials Science and Engineering and

Laboratory for Research on the Structure of Matter

University of Pennsylvania

Philadelphia, PA 19104

Abstract

The crystal-momentum dependence of the line shapes appearing in photoexcitation from the valence levels of ordered S on Fe(100) has been measured using angle resolved photoelectron spectroscopy. The variation in line shape with \vec{k} is a consequence of the hybridization of the energy bands of the 2-D overlayer with the 3-D solid. This observation, predicted theoretically by Liebsch, yields detailed information about the bonding interactions of ordered chemisorption systems.

PACS numbers: 73.20.Hb, 79.60.Gs

UNIVERSITY OF PENNSYLVANIA
LIBRARY

Availability Codes	
Dist	Avail and/or Special
A1	

The physical origin of line shapes in the photoexcitation process has been a subject of great interest for many decades. This has led to a number of descriptive models. The simplest mechanism for line broadening in photoelectron spectroscopy is the hole lifetime. A hole created by the absorption of a photon can decay, primarily by the Auger process or by photon emission, leading to a Lorentzian line shape.¹ Additional broadening may occur by the coupling of the photoexcitation process to many-electron excitations (electron-hole pair creation) or phonons of the system. The phonon coupling leads to a temperature dependent Gaussian broadening,^{2,3} while creation of electron-hole pairs in a metal produces an asymmetric line shape as a result of the Mahan, Nozieres and DeDominicis⁴ x-ray edge singularity.

In a recent angle-resolved experiment, Kevan has argued that the observed anomalous $k_{||}$ -dependence ($k_{||}$ = crystal momentum parallel to the sample surface) of the line width of the sp surface state on Cu(111) is a ground state phenomenon resulting from surface defects.⁵ We report in this letter the first observation of momentum dependent line widths for photoemission from the valence levels of an ordered 2-dimensional overlayer on a single crystal substrate. The chemisorption system used for this work was c(2x2)S/Fe(100). The line shape effect we see is a one-electron property, originating in the interaction (hybridization) of the overlayer states with the substrate continuum. This effect, which is characteristic of the ideal ($k_{||}$ -conserving)

surface, has been predicted by Liebsch⁶, and is reminiscent of the line widths of intrinsic surface states vs. surface resonances. It will be demonstrated that momentum dependent line width data for an ordered overlayer yields new information concerning the bonding of adsorbate atoms to a surface.

Liebsch has calculated momentum dependent line widths for the system $c(2 \times 2)$ oxygen on $Ni(100)$.⁶ Figure 1 is taken from his paper and will be utilized to illustrate the physics of the phenomena described in this article. For the sake of brevity we will describe only the results of the calculation and our data along the $\bar{\Gamma}\bar{M}$ line in the surface Brillouin zone (SBZ-see Fig. 1b). Panel (c) of Figure 1 shows the calculated energy bands for the isolated oxygen layer with the spacing appropriate to the $c(2 \times 2)O$ on $Ni(100)$ geometry. The levels formed from the p_z and p_x oxygen wave functions are even with respect to the mirror plane defined by the $\bar{\Gamma}\bar{M}$ direction and the z -axis (perpendicular to the plane), while the p_y levels are odd. Panel (d) illustrates the effect of the Ni substrate on the oxygen levels. The shaded regions of this panel indicate the projections of the 3-D Ni energy bands onto the SBZ. Only the even Ni bands are shown, since the odd symmetry bands lie energetically above the smallest oxygen binding energy of Fig. 1d. The cross-hatched regions indicate the broadening of the O 2p-derived levels along the bands upon chemisorption. The basic results can be summarized:

- 1) The bonding for this system is principally through the $O(p_z)$ -Ni(sp) interaction.
- 2) The position and shape of the oxygen-derived bands have changed appreciably due to interaction with the Ni substrate.
- 3) The even p_z and p_x derived oxygen bands hybridize because the inversion symmetry of the O layer is broken by the presence of the Ni substrate.
- 4) The oxygen derived levels exhibit an appreciable width when they overlap the projection of the bulk bands, due to hybridization with the Ni levels. The width of each level at a given $k_{||}$ is then a direct measure of the hybridization of the adsorbate band with the substrate band. Note that $k_{||}$ is a good quantum number for both the 2-D and 3-D systems, so only levels with the same energy and $k_{||}$ can mix. For example, the p_z level at $\bar{\Gamma}$ has ~ 0.5 eV width. In contrast, the p_z level at \bar{M} and the odd-symmetry p_y levels for all $k_{||}$ lie outside of the projection of the bulk bands and are consequently discrete.

The measurements described here were made at the Tantalus I storage ring of the University of Wisconsin using the University of Pennsylvania beam line. The procedure for these types of experiments has been described in detail in other articles.⁷ The sample was prepared by segregating sulfur to the surface of a clean Fe(100) single crystal in order to form a c(2x2) LEED pattern.⁸ Previous LEED measurements have determined that the S sits in a four-fold site at 1.1 Å above the Fe(100) plane.⁹

Smooth backgrounds were subtracted from the measured photoemission spectra, and the S features were deconvoluted by Gaussian fitting. (The observed S-peaks were well described by Gaussians, which were chosen for their computational simplicity. Lorentzian fitting does not alter the basic results of this study.) Corrections were made for lineshape broadening effects due to the finite angular acceptance of the electron analyzer. The recorded widths are believed to be accurate within $\pm 10\%$.

Figure 2 shows the measured dispersion and level width at each $k_{||}$ of the $3p_y$ -derived S band, whose levels are of odd symmetry with respect to the $\bar{\Gamma}\bar{M}$ mirror plane. The odd symmetry Fe bands lie energetically above the band of Fig. 2a and therefore can not hybridize with it. Thus the measured linewidths must be a consequence of lifetime broadening and any additional many-particle effects associated with the photoexcitation process. Figure 2b plots the full width at half maximum (FWHM) of the levels along this band as a function of $k_{||}$. The measured width is $1.0 \text{ eV} \pm 0.15$, independent of energy or $k_{||}$.

Figure 3a shows the measured dispersion of the even S bands from the zone center at $\bar{\Gamma}$ out to the second zone center. (The shaded region shows the projection of the even Fe bands.) It is easy to see the hybridization of the p_x and p_z bands along $\bar{\Gamma}\bar{M}$. The top band starts out with a pure p_x -derived level at $\bar{\Gamma}$ and disperses downward until it mixes with the p_z band. Thus, the top band level at \bar{M} is pure p_z in character.⁶ Figures 3b and c plot

the hybridization contribution to the widths after removal of a constant width of 0.85 eV. This width was assumed to be constant for all $k_{||}$ based on the odd-band results of Fig. 2b. The value of 0.85 eV was selected to yield zero hybridization broadening at \bar{M} , where the narrowest levels (even and odd) are observed. The shading in these figures indicates where the bands lie on top of the projected Fe sp-bands, as determined from Fig. 3a. The functional behavior of the linewidths can be qualitatively explained using Liebsch's model:

1) Figure 3b shows the hybridization width for the top band of Fig. 3a. At $\bar{\Gamma}$, the p_x -derived level is not broadened since it can not mix with the substrate s-p band for symmetry reasons.⁶ Upon leaving $\bar{\Gamma}$, this level no longer has the full p_x symmetry and broadens by mixing with the Fe sp-levels. Once it has crossed out of the projected Fe bands, however, the level returns at \bar{M} to the intrinsic width seen for the odd-band levels.

2) Figure 3c shows the hybridization widths for the levels along the bottom band of Fig. 3a. At $\bar{\Gamma}$, the width is very large (~ 2 eV) due to the large hybridization of the S p_z level with the Fe sp-levels of the same $k_{||}$. The width becomes gradually smaller in proceeding across the SBZ as more p_x character is introduced, and less hybridization with the bulk levels occurs. Finally, the band crosses out of the projected Fe bands and the levels show almost no hybridization broadening at \bar{M} , indicating that lifetime broadening is responsible for the widths near this

point.

When the above results for $c(2 \times 2)S/Fe(100)$ are compared with a previous photoemission study of $c(2 \times 2)S/Ni(100)$,¹⁰ and the theoretical calculation for $c(2 \times 2)O/Ni(100)$,⁶ it can be concluded that the $S(3p_z)-Fe(sp)$ interaction is much more pronounced in this system. Thus the observation of $k_{||}$ -dependent linewidths can provide important information about the nature of the chemisorption bond.

There are two observations that we made about this system that were not anticipated by Liebsch's calculation.⁶ The first is that the hybridization line widths shown in Figure 3 continue to vary with $k_{||}$ even after the S level has moved out of the bulk band projections, qualitatively similar to Kevan's observation for the surface state on $Cu(111)$.⁵ The effect we observe is too large to attribute to $k_{||}$ smearing due to surface defects, however. This effect could result from the coupling of the hole lifetime width with the $k_{||}$ -resolved hybridization width. In an independent particle picture similar to the one used by Liebsch⁶, the S levels are discrete when they lie outside of the projection of the bulk bands. In Fig. 2, however, there is ~ 1 eV width in these regions due to the hole lifetime. This finite lifetime for the state characterized by $E(k_{||})$ implies an uncertainty in $k_{||}$ and E allowing bulk 3-D states to hybridize with surface 2-D states even when they appear to be separated in $E-k_{||}$ space.

The second observation is depicted in Fig. 4, where we show

two measured energy distributions. The collection geometry is normal to the crystal, ($\bar{\Gamma}$ in the SBZ), and the photon energy and polarization of the incident light have been varied. The dashed curves are for nearly s-polarized light where only the $p_x p_y$ -levels can be excited, while the solid curves are for p-polarized light where all levels can be excited. As the photon energy is changed, the energy position and width of the $p_x p_y$ level is unchanged as it should be for a two dimensional system.⁷ In contrast, the p_z level moves in energy and its width increases appreciably as $h\nu$ is changed from 25 to 30 eV. At first glance, this observation seems to violate the contention that we are looking at a 2-D band structure where the levels are described by $E(k_{||})$. The origin of this shift in the p_z level with photon energy is the large hybridization of the S p_z with the Fe s-p bands. This feature in the photoemission spectra contains large admixtures of S p_z and Fe s and p wave function character, with each having a slightly different local density of states. As the photon energy is changed the cross-sectional variation of each component is different resulting in apparent changes in the line shape and position.

This work was supported by ONR. The University of Pennsylvania beam line is supported by NSF through MRL Contract No. DMR-8216718. We would like to thank the staff of the Synchrotron Radiation Center at the University of Wisconsin for their assistance, and Prof. P. Soven for his helpful comments.

References

1. John C. Fuggle and Santos F. Alvarado, Phys. Rev. A22, 1615 (1980) have measured the life time of core holes in approximately 25 elements and compared the lifetimes to theoretical predictions.
2. G.K. Wertheim and P.H. Citrin, in Photoemission in Solids I, edited by M. Cardona and L. Ley (Springer, Berlin, 1978), p. 197.
3. W.R. Salaneck, C.B. Duke, W. Eberhardt, E.W. Plummer, and H.J. Freund, Phys. Rev. Lett. 45, 280 (1980).
4. G.D. Mahan, Phys. Rev. 163, 612 (1967); P. Nozieres and C.T. DeDominicis, Phys. Rev. 178, 1097 (1969).
5. S.D. Kevan, Phys. Rev. Lett. 50, 526 (1983).
6. A. Liebsch, Phys. Rev. B17, 1653 (1978).
7. E.W. Plummer and W. Eberhardt, Adv. in Chem. Phys. 49 (1982).
8. R.A. DiDio, Ph.D. Thesis, Univ. of Pennsylvania, 1983 (unpublished).
9. K.O. Legg, F. Jona, D.W. Jepsen, and P.M. Marcus, Surf. Sci. 66, 25 (1979).
10. E.W. Plummer, B. Tonner, N. Holzwarth and A. Liebsch, Phys. Rev. B21, 4306 (1980).

Figure Captions


Fig. 1 (a) Real space unit cell for $c(2 \times 2)O$ on $Ni(100)$ system (X's are O atoms), (b) reciprocal space Brillouin zone for clean $Ni(100)$ (Solid lines) and $c(2 \times 2)O$; (c) energy bands for isolated oxygen layer; and (d) energy levels for $c(2 \times 2)O$ chemisorbed on $Ni(100)$. The shaded regions indicate the projection of the bulk bands.

Fig. 2 Dispersion of the odd symmetric p_y band. (b) Line-width (FWHM) of the levels along the p_y band.


Fig. 3 (a) Dispersion of the two even S bands. The shaded region is the projection of the bulk bands. (b) Hybridization widths along the upper band in (a). (c) Hybridization widths along the lower band in (a).

Fig. 4 Normal emission spectra of $c(2 \times 2)S$ on $Fe(100)$ illustrating the k_{\parallel} -dependence of the S $3p_z$ lineshape at $k_{\parallel} = 0$. (Dashed and solid lines indicate s- and p-polarized incident radiation, respectively.)


(a) SURFACE UNIT CELL




(b) SURFACE BRILLOUIN ZONES




(c) OXYGEN BANDS




(d) CHEMISORBED OXYGEN LAYER




(FIG. 1)



(FIG. 2)



(FIG. 3)



(FIG. 4)

